

## DISTANT HEALING OF SURGICAL WOUNDS: AN EXPLORATORY STUDY

Marilyn Schlitz, PhD,<sup>1</sup> Harriet W. Hopf, MD,<sup>2</sup> Loren Eskenazi, MD, FACS,<sup>3</sup> Cassandra Vieten, PhD,<sup>1</sup> and Dean Radin, PhD<sup>1, #</sup>

**Background:** Distant healing intention (DHI) is one of the most common complementary and alternative medicine (CAM) healing modalities, but clinical trials to date have provided ambivalent support for its efficacy. One possible reason is that DHI effects may involve variables that are sensitive to unknown, uncontrolled, or uncontrollable factors.

**Objective:** To examine 2 of those potential variables—expectation and belief—we explored the effects of DHI on objective and psychosocial measures associated with surgical wounds in 72 women undergoing plastic surgery.

**Design:** Participants were randomly assigned to 1 of 3 groups: blinded and receiving DHI (DH), blinded and not receiving DHI (control), and knowing that they were receiving DHI (expectancy). Outcome measures included collagen deposition in a surrogate wound and several self-report measures. DHI was provided by experienced distant healers. No differences in the main measures were observed across the three groups.

**Results:** Participants' previous belief in the efficacy of DHI was negatively correlated with the status of their mental health at the end of the study ( $P = .04$ , 2-tailed), and healers' perceptions of the quality of their subjective "contact" with the participants were negatively correlated both with change in mood ( $P = .001$ ) and with collagen deposition ( $P = .04$ ). A post-hoc analysis found that among participants assigned to receive DHI under blinded conditions, those undergoing reconstructive surgery after breast cancer treatment reported significantly better change in mood than those who were undergoing purely elective cosmetic surgery ( $P = .004$ ).

**Conclusion:** If future DHI experiments confirm the post-hoc observations, then some of the ambiguity observed in earlier DHI studies may be attributable to interactions among participants' and healers' beliefs, their expectations, and their motivations.

(*Explore* 2012; 8:223-230. © 2012 Elsevier Inc. All rights reserved.)

## INTRODUCTION

Distant healing intention (DHI) may be defined as "a compassionate mental act intended to improve the health and well-being of another person at a distance."<sup>1</sup> Terms used to describe DHI interventions include intercessory prayer, spiritual healing, nondirected prayer, intentionality, energy healing, shamanic healing, nonlocal healing, noncontact therapeutic touch, and Reiki. Each of these methods involves a distinct theoretical, theological, cultural, or pragmatic approach toward healing through the application of one person's intention towards another.

DHI, especially in the form of prayer, is one of the most commonly used complementary and alternative medicine (CAM) healing modalities. A government survey of adult Americans by the U.S. National Center for Health Statistics showed that of the top 10 CAM practices, the first was prayer for oneself and the second was prayer for others.<sup>2</sup> Among social workers, a survey found that 28% of more than 2000 respondents had engaged in verbal prayer with their clients, whereas 57% privately prayed for their clients.<sup>3</sup> In a survey among 1900 cancer survivors, 62% reportedly prayed for their own health, 39% had others pray for their health, and 15% participated in group prayer.<sup>4</sup> And, on the basis of general population surveys from 2002 to 2008, the use of prayer for health concerns has continually increased in the United States after taking into account demographic, socioeconomic status, health status, and lifestyle behaviors.<sup>5</sup>

Although the popularity of DHI is well established, its efficacy is uncertain. In a meta-analysis, Hodge<sup>6</sup> reviewed 17 randomized controlled studies of DHI and concluded that the outcomes produced small but significant effects for intercessory prayer under both random and fixed effect models, with and without inclusion of one controversial study.<sup>7</sup> Masters et al<sup>8</sup> focused on 14 studies involving only intercessory prayer and found a positive but nonsignificant outcome. In an updated meta-analysis with one additional study, Masters et al<sup>9</sup> again found a positive, nonsignificant outcome. That data also showed that the DHI effect size, although small, was nearly 15 times larger for unhealthy subjects than for healthy controls, suggesting that need or motivation may play a role in DHI efficacy. Nevertheless, because the overall results were not robust, Masters et al recommended that further resources not be allocated

1 Institute of Noetic Sciences, Petaluma, CA

2 Department of Anesthesia and Perioperative Care, University of California, San Francisco, San Francisco, CA

3 Plastic Surgery Associates, San Francisco, CA

Current address of Harriet W. Hopf, MD: Department of Anesthesiology, University of Utah, School of Medicine, Salt Lake City, UT 84132. This publication was made possible by grant number R21AT001437 from the National Center for Complementary and Alternative Medicine (NCCAM) at the National Institutes of Health. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of NCCAM. None of the authors have any financial relationships or other relationships that may pose a conflict of interest with the aims or outcomes of the present study.

# Corresponding Author. Address:

Institute of Noetic Sciences, 625 Second Street, Suite 200, Petaluma, CA 94952-9524

e-mail: dean@noetic.org

---

for this type of research. The most recent meta-analysis, a Cochrane systematic review, Roberts et al<sup>10</sup> found fewer deaths in an intercessory prayer condition as compared with standard care control, results that were again significant for high risk and thus highly motivated subjects ( $P < .00001$ ). Their conclusion was similar to that of Masters et al: "The evidence presented so far is interesting enough to support further study. However, if resources were available for such a trial, we would probably use them elsewhere" (p. 24).

Laboratory evidence for the possibility of DHI efficacy specifically on the healing of wounds was first published by Grad et al,<sup>11</sup> who studied the effects of a noted healer, Oskar Estebany, on surgical wounds in 48 mice as compared with a nonhealed control group. In a study that was later successfully replicated, mice exposed to the healer showed significantly enhanced wound healing, suggesting a DHI effect not attributable to placebo effects. In a more recent study that also involved mice but with DHI directed towards healing of injected mammary cancer rather than surgical wounds, Bengston<sup>12</sup> also found significantly enhanced healing.

Overall, clinical trials and laboratory studies suggest that there might be some efficacy for DHI but lack of a clear consensus and uncertainty about the key underlying variables has led most analysts to recommend that research funding be directed towards more tractable problems. For pragmatic reasons, such advice is not unreasonable. However, if DHI effects are genuine to any extent, even if they manifest only as a small magnitude, high variance phenomenon, then the scientific and medical consequences are profound and additional research is justified. In particular, exploring the extent to which effects of DHI may be attributable to or modulated by patient expectations and resulting placebo effects, or how patient beliefs or motivations might interact with DHI, offers the potential to shed light on the mind-body healing process in general.

In this study, we chose to explore the effectiveness of DHI on the healing of surgical wounds. For an objective outcome measure we used the rate of subcutaneous collagen deposition. To address other possible outcomes of DHI, we also administered a set of self-report measures of mental and physical health.

In the real world, people often know that they are being treated with DHI, so it can be difficult to determine what proportion of the reported benefit may be due to DHI and what may be attributed to expectancy or placebo. To explore the role of expectation in DHI, we designed a 3-arm study: a blinded group receiving DHI, a blinded group not receiving DHI, and an unblinded, placebo-enhanced group receiving DHI. This allowed us to study the effects of subjects' expectations by varying whether the subject knew that DHI efforts were being made on their behalf. The primary research questions were as follows: (1) Do participants' expectations about whether they are the target of DHI influence collagen deposition and is this correlated with their subjective measures of well being? (2) Do participants' beliefs about DHI affect collagen deposition or subjective measures of well being?

## METHODS

### Procedure

The Institutional Review Boards of the California Pacific Medical Center and St Mary's Medical Center, both in San Fran-

cisco, approved this study and informed consent was obtained from all participants. The study involved women undergoing elective surgery, some for reconstruction after breast cancer surgery and some for cosmetic reasons. They were randomized into 3 groups:

1. A control group knew whether they were being randomly assigned to receive DHI, and in this condition they did not receive it.
2. A distant healing (DH) group knew whether they were being randomly assigned to receive DHI, and in this condition they did receive a 20-minute or longer DHI session per day for 8 days.
3. An expectancy group knew that they were receiving DHI in a 20-minute or longer DHI session each day for 8 days, and they were reminded of this daily by a telephone call.

All participants, clinical staff, and data analysts were blinded to group assignments, and for the expectancy group, the research staff and data analysts were blinded to group assignment.

During the study period, all women entering the Plastic Surgery Associates facility (San Francisco, CA) who were eligible for the study were contacted by mail before their preoperative visit. A recruitment letter described the study and provided contact information if they had questions about the study before they arrived at the office. The recruitment letter specified that this study concerned "distant healing, or someone holding the intention for your healing in a remote location." Participants who expressed interest had an enrollment interview on site at Plastic Surgery Associates approximately 2 weeks before surgery. Study participants also filled out initial questionnaires and were photographed. After surgery, participants engaged in their daily activities with no special requirements from the study. They were contacted for a three-month follow-up interview administered via mail. Contacts between participants and study personnel were kept to a minimum.

### Assessments

Using an investigator-developed Personal Data Questionnaire (PDQ), we collected demographic information, baseline health behaviors including nutritional intake and exercise, use of alternative therapy interventions including other forms of DHI, and degree of social support. It also contained a 5-point Likert scale to assess the subject's degree of belief in the effectiveness of DHI as a healing modality.

**IMPRA implants.** Hydroxyproline (collagen) deposition was measured at the 8-day postsurgical assessment point by the use of 2 expanded polytetrafluoroethylene implants (International Polymer Engineering, Inc, Tempe, AZ, trade name IMPR), positioned subcutaneously following a standard method described in the wound healing literature. A total of 13 IMPRA samples were stored at  $-70^{\circ}\text{F}$  until the time of assay. Quantification of collagen deposition was made

---

through chromatographic analysis for hydroxyproline, using micrograms per centimeter units. The assay was performed according to the methods of Lindblad and Diegelmann.<sup>13</sup>

The *Short Form Health Survey* (SF-36) is a 36-item questionnaire that generates a health-related quality of life profile consisting of 8 scales and 2 summary measures: a Physical Composite Score (PCS) and a Mental Composite Score (MCS).<sup>14</sup> Used in more than a thousand published studies, the original SF-36 has proven to be useful in monitoring general and specific populations, comparing the burden of different diseases, differentiating health benefits produced by different treatments, and screening individual participants.

The Profile of Mood States (POMS) is a 65-item measure of level of mood disturbance; it takes approximately 3 to 5 minutes to complete.<sup>15</sup> The scale provides a total score and 6 dimensions of mood, including tension, depression, anger, vitality, fatigue, and confusion. It was first assembled and tested for validity and reliability in 1955. Internal consistency reliabilities based on a sample of female outpatients range from 0.84 (confusion-bewilderment) to 0.95 (depression-dejection).

The *Rosenberg Self-Esteem Scale* is a brief, single-scale validated measure of self-esteem.<sup>16</sup>

The *Impact of Surgery Scale* is a 5-point Likert investigator-developed instrument designed to capture the subject's overall perception of the impact of plastic surgery on their lives in retrospect. It was administered via mail 3 months after the surgery.

### **Surgical Procedure**

All surgical procedures were performed by one of the authors (L.E.), a licensed plastic surgeon, at California Pacific Medical Center or at St Mary's Hospital (San Francisco, CA). At the end of the surgery, while the subject was still under anesthesia, a small wound was made with a number 11 blade at the planned entry and exit sites of the IMPRA collagen collection tube. Two 1-mm × 7-cm sections of IMPRA tubing were implanted. Two tubes were implanted to increase sample volume because 7 cm is the limit of tube length that can be easily implanted subcutaneously. The same entry site was used for both implants, and the ends of each implant protruded from a separate exit site. All IMPRA implants were placed in a standardized location in the lower portion of the abdomen, just above the pubis.<sup>17</sup> This standardized surrogate site was chosen because it represents the same tissue compartment as the actual wound, but is far enough from the wound to alleviate concerns for contamination. Participants were informed that placement of the tubes may leave a small (1-2 mm) scar. Because many of the women in the study were having cosmetic and/or reconstructive surgery, and would thus likely be concerned about such a scar near their surgical wound, we chose a site where a scar would not be an issue. Using a surgical needle, the surgeon pulled the IMPRA through, just under the skin, until the distal tip disappeared under the skin and only a few millimeters protruded proximally. They then dressed the wound with a thin, semipermeable, transparent film dressing (Tegaderm; 3M, St Paul, MN).

On the eighth day after surgery, the subject returned to the Plastic Surgery Associates office to have the surgical sutures removed. The surgeon also removed the IMPRA tubes from the

implant site. A topical anesthetic (EMLA cream; AstraZeneca LP, Wilmington, DE) was used on the exit site to minimize the pain of removal. The dry, exposed end of the IMPRA implant was cut off and the remainder placed in separate test tubes, then placed on ice, coded to match the subject code, and stored in a -70°F freezer within 2 hours of removal. Samples were then delivered on ice to an analytical laboratory at the Wound Healing Laboratory at the University of California, San Francisco, where they were stored in a -20°F freezer until analysis. To reduce assay variability, an analysis was performed simultaneously on all samples after subject enrollment was complete. The hydroxyproline content for the 2 implants removed from each subject was measured separately. Values were then averaged and the single average value used for analysis, as in previous studies.<sup>18</sup> Ingrowth into the implants is somewhat heterogeneous and this method provides the most accurate representation of the healing capacity of the subject.<sup>18,19</sup>

If the subject did not have a follow-up appointment scheduled on the eighth day after surgery and did not wish to come in to the Plastic Surgery Associates office for an additional appointment (some participants lived far enough away that this constituted a hardship, particularly during recovery from surgery), a nurse trained in IMPRA implant removal removed the implants at the subject's home.

### **DHI Intervention**

The study was designed to explore the use of DHI as it is commonly practiced by professional healers, ie, without providing precise prescriptions instructing the healers what to do. Participants in the treatment conditions received healing effort from DHI practitioners who were asked to "Direct an intention for the health and well-being of the subject, for at least 20 minutes per day for eight days." When a new subject entered the study, the DHI practitioner assigned to work on the subject was provided (by overnight mail delivery) with a subject information sheet including a 4- by 6-inch (10.2-cm by 15.2-cm) color photograph, the subject's first name, and the date and approximate time of surgery. Each sheet was placed in a sealed envelope with instructions not to open it until the date the DHI practitioner was to begin treatment (ie, the day of surgery). The DHI practitioner's location and distance from the subject was unrestricted. Each subject received DHI by one practitioner, and each practitioner was randomized to direct DHI to two participants (one at a time, separated by at least one month). This design was intended to help distribute possible variations in DHI practitioners' healing abilities as uniformly as possible across the subject pool, and also to give DHI practitioners adequate respite.

Each DHI practitioner was asked to commit 20 minutes of their time per day per patient, plus time to complete the healer daily log. DHI practitioners were asked to treat their subject for 8 days in a row and to fill out a 2-page, 8-question log to record time, location and duration of each day's healing effort, as well as any comments, images or impressions that they wished to report. The log also contained a checklist that documented the use of any of 25 different DHI strategies, such as "sending light," "praying to a higher power," "visualizing the subject well," etc., as well as any intuitive diagnostic efforts made. They were also asked

to report to what degree they felt “affinity for, resonance with, or connection to” the subject, and how effective they felt the healing session was. DHI practitioners received a reminder phone call the day before they were scheduled to begin their 8 days of DHI, a thank you phone call and letter at the close of each 8 days of DHI, and a small gift, but they were not paid for their participation.

### Data Analysis

Participants were compared on baseline demographics to assess similarity across groups. Baseline variables showing significant between-group differences were controlled in subsequent analyses via analysis of covariance (ANCOVA). Spearman rank-order correlations were used to examine relationships between expectancy ratings and healing outcomes. All reported *P*-values are 2-tailed. This study was explicitly exploratory. It was intended to examine possible trends and to establish effect sizes for future confirmatory studies.

Four outcome measures were used: natural log (to adjust for a positively skewed distribution) of collagen deposition, change in POMS normalized mood scores from baseline to end of the 8-day DHI period, SF-36 MCS at the end of the DHI period, and SF-36 PCS at the end of the DHI period.

### Participant Selection

**Inclusion criteria.** Inclusion criteria were as follows: (1) women undergoing plastic surgery; and (2) American Society of Anesthesiologists (ASA) Physical Status Criteria I or II (ie, ASA I = health excellent with no systemic disease; no limitations on activity; no danger of death; excluded: persons at

extremes of age; ASA II = health: disease of one body system; status of underlying disease is well-controlled; no limitations on activity). Participants were compensated \$100 in the form of a check upon completion of all study requirements.

**Exclusion criteria.** Exclusion criteria were as follows: (1) non-English speaking; (2) inability or unwillingness to fill out questionnaires; (3) being a smoker; (4) morbid obesity; (5) circulatory inadequacies; (6) nutritional deficits as evidenced by neutropenia or hypoalbuminemia; (7) history of radiation to the site of IMPRA implant, or any radiation therapy within the past 3 months, (8) current stage 3 or 4 breast cancer; (9) duration of surgery anticipated to be < 2 hours.

**Healer inclusion criteria.** Healers identified for this study had: (1) a minimum of 5 years of experience as a professional DHI practitioner, meaning that they regularly used DHI for treatment of participants or clients not otherwise known to the DHI practitioner; (2) had practiced healing at a distance on at least 30 previous patients; and (3) had experience working with wound healing. Healers were recruited from a database of more than 100 healers maintained by California Pacific Medical Center Research Institute, all of whom had been previously evaluated and met or exceeded these criteria. Applicants filled out a Healer Information Form which provided verification of inclusion criteria and additional descriptive information. DHI practitioners were located in North America within a range of 3 to 3000 miles away. Practitioners were experienced in a variety of DHI

**Table 1.** Subject Demographics

|                       |       | Ethnicity         |    | Education           |    | Reason for Surgery |    |
|-----------------------|-------|-------------------|----|---------------------|----|--------------------|----|
| Control group         |       | European-American | 19 | High school diploma | 2  | Cancer             | 10 |
| Participants          | 24    | Other             | 4  | Technical School    | 1  | Cosmetic           | 14 |
| Average age           | 44    | Nonreporting      | 1  | Some college        | 1  |                    |    |
| Age range             | 28-63 |                   |    | College degree      | 13 |                    |    |
|                       |       |                   |    | Graduate degree     | 7  |                    |    |
|                       |       |                   |    |                     |    | Total              | 24 |
| Distant healing group |       | European-American | 21 | High school diploma | 1  | Cancer             | 11 |
| Participants          | 23    | Other             | 0  | Technical School    | 0  | Cosmetic           | 12 |
| Average age           | 49    | Nonreporting      | 2  | Some college        | 5  |                    |    |
| Age range             | 28-63 |                   |    | College degree      | 8  |                    |    |
|                       |       |                   |    | Graduate degree     | 7  |                    |    |
|                       |       |                   |    | Nonreporting        | 2  |                    |    |
|                       |       |                   |    |                     |    | Total              | 23 |
| Expectancy group      |       | European-American | 20 | High school diploma | 2  | Cancer             | 7  |
| Participants          | 25    | Other             | 3  | Technical School    | 1  | Cosmetic           | 17 |
| Average age           | 48    | Nonreporting      | 2  | Some college        | 7  | Both               | 1  |
| Age range             | 32-67 |                   |    | College degree      | 10 |                    |    |
|                       |       |                   |    | Graduate degree     | 3  |                    |    |
|                       |       |                   |    | Non-reporting       | 2  |                    |    |
|                       |       |                   |    |                     |    | Total              | 25 |

modalities, including visualization, prayer, energy work, and shamanic healing.


## RESULTS

The study involved 40 healers who collectively directed DHI towards 72 women, aged 28 to 63. Table 1 lists subject demographics and the reason for surgery. After receiving surgery, no participants dropped out of the study because they had to visit the surgeon's office or see a nurse to have the IMPRA implant removed, and at that time they were all available to fill out the 8-day postsurgery questionnaires. However, 17 participants failed to return the mailed 3-month follow-up questionnaire (24%). The drop-outs were spread approximately evenly among the 3 groups (control: 29%; DH: 17%; expectancy: 24%). Of 48 participants who were assigned a healer (23 in the DH group, 25 in the expectancy group) there were 2 instances in which the assigned healer did not complete the daily logs (one in the DH group, one in the expectancy group). For the remaining 46 sessions, all daily logs were completed, and many healers provided extensive notes on the sessions.


### Aim 1: Did Expectations About DHI Influence Collagen Deposition or Subjective Measures of Well Being?

Examination of demographic and other variables revealed one significant difference among the groups: hours of surgery (Figure 1). Surgeries in the expectancy group participants lasted nearly an hour longer on average than the control group, and about 50 minutes longer than the DH group ( $F(2,64) = 5.22, P = .008$ ). Participants were randomly assigned and the surgical staff, and research team were blinded to assignment at the time of surgery, so it appears that randomization failed to equally distribute surgical times in this study. As a result, this factor was controlled for by being entered as a covariate in all subsequent analyses.

**Collagen deposition.** There was no significant difference in mean collagen deposition among the three groups ( $F(2,62) = .79, P = .46$ , see Figure 2).



**Figure 1.** Means and one standard error of the mean bars for hours of surgery in the control, DH, and expectation groups.




**Figure 2.** Means and one standard error of the mean error bars for natural log of collagen deposition for the control group (blinded, not receiving DHI), the DHgroup (blinded, receiving DHI), and the expectation group (unblinded, receiving DHI).


**Subjective measures.** There were no mean changes in mood among the 3 groups ( $F[2, 48] = 2.54, P = .09$ , Figure 3). There were also no significant mean differences in the SF-36 PCS (physical composite) scores ( $F[2, 56] = 2.08, P = .13$ , see Figure 4) or the SF-36 MCS (mental composite) scores ( $F[2, 56] = 2.41, P = .10$ ). However, a Fisher LSD (least significant difference) post-hoc pairwise comparison did suggest that mood change in the DHI group was more positive than in the Control group ( $P = .02$ , 2-tailed).

### Aim 2: Did Participants' Beliefs About DHI Affect Collagen Deposition or Subjective Measures of Well Being?

To assess this aim, we evaluated Spearman rank order correlations across all participants (ie, without regard to group assignment) for the subjective measures (change in mood, and SF-36



**Figure 3.** Means and one standard error bars for change in mood (POMS) scores from day of surgery to end of DHI period 8 days later.



**Figure 4.** Means and one standard error bars for SF-36 PCS and MCS on the last day of the 8-day DHI period.

PCS and SF-36 MCS on the last day of DHI) and objective measures (natural log of collagen deposition) versus their (1) prior belief in the efficacy of DHI, (2) belief that they were assigned to a group that would receive DHI, (3) belief that DHI was actually taking place, and (4) the degree to which the healers felt “connected” to the participants.

Participants’ previous belief in DHI was significantly negatively associated with their mental state at the end of the study period (SF-36 MCS, Spearman rho =  $-.27$ ,  $P = .04$ ). Similarly, the degree to which participants’ felt that DHI was taking place was weakly and negatively associated with collagen deposition (Spearman rho =  $-.31$ ,  $P = .06$ ).

To examine these results in more detail, we partitioned the data into the 3 groups. The Expectancy group’s results were consistent with the combined analysis, ie the more that participants believed that they were receiving DHI, the worse their mental state was at the end of the study period (SF-36 MCS, Spearman rho =  $-.50$ ,  $P = .02$ ). Also, the more that they believed that DHI was actually taking place, somewhat less collagen was deposited (Spearman rho =  $-.38$ ,  $P = .06$ ). Furthermore, the more that healers reported a sense of connectedness with their patients, the more negative was the participants’ change in mood (Spearman rho =  $-.56$ ,  $P = .03$ ) and the less collagen was deposited (Spearman rho =  $-.44$ ,  $P = .07$ ). In the DH group, the healers’ sense of connection with the participants, and the participants’ change in mood, were again negative (Spearman rho =  $-.62$ ,  $P = .01$ ). The control group showed no significant relationships or trends.

#### Post-hoc Analysis

In a series of post-hoc analyses, we examined other variables that may have affected the results. To further explore the correlations observed in the expectancy and DH groups, we investigated whether those outcomes may have been influenced by the participants’ motivation for electing surgery. Participants who underwent surgery for cosmetic purposes accounted for the majority of study drop-outs (eg, more cosmetic surgery vs. reconstruction patients failed to complete follow-up measures). Furthermore,


the expectancy group contained a greater proportion of cosmetic surgery patients than the control or DH groups. To explore whether differences in the purpose of surgery might have differently affected the two group’s receptiveness to DHI,<sup>20</sup> we partitioned participants into those receiving reconstructive surgery after treatment for breast cancer ( $n = 28$ ) versus those who had surgery for cosmetic purposes ( $n = 44$ ).

There was a significant interaction between change in mood and motivation for surgery (analysis of covariance group  $\times$  diagnosis  $\times$  change in POMS:  $F(2,45) = 3.56$ ,  $P = .04$ , Figure 5). A post-hoc Fisher LSD pair-wise comparison showed that the reconstructive surgery participants in the DH group experienced significant improvements in mood as compared with the cosmetic surgery participants in any other condition: cosmetic control ( $P = .002$ ), cosmetic DH ( $P = .004$ ), cosmetic expectancy ( $P = .01$ ), reconstructive control ( $P = .001$ ), or reconstructive expectancy ( $P = .06$ ). No significant interactions were observed between the purpose of surgery versus PCS or MCS scores, or for collagen deposition.

Finally, healers’ self-rated sense of “connectedness” with their participants, based on their subjective impressions recorded on their daily healing logs, was negatively associated with both participants’ change in mood (Spearman rho =  $-0.57$ ,  $P = .001$ ) and with the amount of collagen deposition measured (Spearman rho =  $-0.30$ ,  $P = .04$ ).

#### DISCUSSION

DHI is one of the most widely used CAM practices and undoubtedly one of the most controversial. The practice is common because in cases of incurable illness it is often the only course of “treatment” that is available, and for many loved ones it may be the only thing they feel they can offer. It is controversial because, as in the present study, the results of clinical trials are rarely clear-cut, and also because some assume that DHI effects are impossible in principle due to a lack of widely accepted theoretical mechanisms specifying how influences could extend from one person’s mind to a distant person’s body.<sup>21</sup>



**Figure 5.** Interaction between mean change in mood scores and purpose of the surgery, with one standard error of the mean error bars.

Some theologians further consider DHI to be controversial because they assume science is either inadequate or inappropriate for testing DHI interpreted as divine intervention,<sup>9</sup> or worse, that if some prayers were answered and others were not, it would imply that God is immoral.<sup>22</sup> Moreover, some bioethicists are concerned that DHI experiments may be unethical because many assume that such interventions are benign and risk-free.<sup>23</sup> It was in part for this reason that our study did not include a deceptive condition where people receiving DHI were unaware of that possibility.

Concern about possible harm is not without anecdotal support (albeit at the level of folklore).<sup>24</sup> For example, some fundamentalist preachers have been known to openly evoke “imprecatory prayer” to pray for the death of government officials with whom they disagree.<sup>25</sup> There is also some empirical support for negative effects. A study of intercessory prayer by Benson et al<sup>26</sup> found that participants who were told that they were receiving prayer were 14% more likely to experience complications than individuals who were also receiving prayer but uncertain of it, or if they were not receiving prayer. In addition, in an earlier prayer study, Walker et al<sup>27</sup> found that participants’ participation in the prayer study (whether being prayed for or not) delayed the reduction of drinking and knowledge that a friend or family member was praying for them at baseline resulted in increased alcohol consumption at 6 months as compared with those reporting they were unaware of anyone praying for them. This finding suggests that complex interactions may occur between actually receiving prayer or other DHI modalities, and one’s knowledge, beliefs and expectancies about whether one is receiving DHI.

The present study was intended to explore these issues. The results suggested no main effects of DHI on the primary outcome measure of collagen deposition, or on secondary measures of self-reported emotional and physical health after surgery. The only significant difference between the conditions was that length of surgery in the Expectancy group was significantly longer than in the DH or Control groups. Because the DHI intervention did not begin until the day following surgery, barring a retrocausal explanation it seems likely that this difference was due to chance, thus to equalize this variable between the three groups it was used as a covariate in subsequent analyses. The most parsimonious interpretation of the main results is that neither DHI nor expectancy affected the outcome measures. It is possible that other limitations in the study design, including a lack of statistical power, may account for the lack of difference between the 3 conditions.

Post-hoc analyses suggested that concerns about possible negative effects of DHI may deserve a closer look. That is, this study did not provide support for the efficacy of DHI on wound healing *unless* participants’ and healers’ beliefs and expectations are taken into account. When those factors are included, a more complex picture emerges: The more that participants believed in distant healing, and the more they thought that distant healing was actually focused on them, the *worse* they did on both objective and subjective measures. In addition, the better the healers thought that they were doing, again, the *worse* the participants’ outcomes. Similar effects were not observed in the control group, suggesting that

DHI played a role in these outcomes. These findings are reminiscent of the results of the Benson et al.<sup>26</sup>

Additional post-hoc findings suggested that the motivation for having surgery may play an important role in modulating DHI effects. Participants’ change in mood was substantially improved in the DH condition in women receiving reconstructive surgery after breast cancer treatment but not in women receiving cosmetic surgery. Limitations of this finding include different proportions of cosmetic versus reconstructive surgeries in each condition and differences in drop-out rates between cosmetic versus reconstructive surgery participants. However, this finding does suggest that motivation for surgery may be a worthwhile variable for further study.

What might account for the generally negative trend associated with belief in DHI? It is conceivable that high belief in DHI may be associated with less psychological resilience to surgery. It is also possible that high-belief patients expect that other people or forces outside of themselves can exert healing effects, and thus they might feel a lower sense of responsibility for their own healing. Another possibility, based on laboratory studies of the effects of distant intention on the human nervous system,<sup>20,28,29</sup> is that distant intention effects are often correlated with activation of the sympathetic nervous system, ie, a stress response. Increased stress is known to have negative effects on wound healing rates,<sup>30</sup> suggesting that participants who are the target of DHI, especially those who do not have strong motivations to heal, may experience a rise in stress, and as a result a decrease in wound healing rates.

In conclusion, one explanation why meta-analyses are providing ambivalent conclusions about the efficacy of DHI is that distant healing effects do not exist. From that perspective, the occasional positive report of a properly conducted DHI study is best attributed to a statistical false positive or to selective reporting. An alternative explanation is that DHI effects do exist, but the relevant variables that modulate these effects are not well understood and interact in complex ways. Ultimately, future empirical studies or development of new theories will be required to help decide which of these explanations is closer to the truth.

#### Acknowledgments

We thank Raymond Buscemi for his contributions to this study, Dan Moore and Jessica Utts for statistical consultation, and Jenny Mathews, Charlene Farrell, Judith West, and Sue Louiseau for their assistance with the project. We also wish to thank David and Marjorie Griswold of For Goodness Sake for their financial support, and Charlene Farrell for her administrative and editorial support.

#### REFERENCES

1. Slicher FT, Targ E, Moore D, 2nd, Smith HS. A randomized double-blind study of the effect of distant healing in a population with advanced AIDS: report of a small scale study. *West J Med.* 1998;169: 353-363.
2. Barnes BM, Powell-Griner E, McFann K, Nahin RL. Complementary alternative medicine use among adults: United States. *Adv Data.* 2002;2004:1-19.

3. Canda ER, Furman LD. *Spiritual Diversity in Social Work Practice: the Heart of Helping*. New York, NY: Free Press; 1999.
4. Mao JJ, Farrar JT, Xie SX, Bowman MA, Armstrong K. Use of complementary and alternative medicine and prayer among a national sample of cancer survivors compared to other populations without cancer. *Complement Ther Med*. 2007;15:21-29.
5. Wachholtz A, Sambamoorthi U. National trends in prayer use as a coping mechanism for health concerns: changes from 2002 to 2007. *Psychol Relig Spirituality*. 2011;3:67-77.
6. Hodge DR. Social justice and people of faith: a transnational perspective. *Soc Work*. 2007;52:139-148.
7. Cha KY, Wirth DP. Does prayer influence the success of in vitro fertilization-embryo transfer? Report of a masked, randomized trial. *J Reprod Med*. 2001;46:781-787.
8. Masters KS, Spielmanns GI, Goodson JT. Are there demonstrable effects of distant intercessory prayer? A meta-analytic review. *Ann Behav Med*. 2006;32:21-26.
9. Masters KS, Spielmanns GI. Prayer and health: review, meta-analysis, and research agenda. *J Behav Med*. 2007;30:329-338.
10. Roberts L, Ahmed I, Hall S, Davison A. Intercessory prayer for the alleviation of ill health. *Cochrane Database Syst Rev*. 2011:1-48.
11. Grad B, Cadoret R, Paul G. The influence of an unorthodox method of treatment on wound healing in mice. *Int J Parapsychol*. 1961;3:5-24.
12. Bengston W. *The Energy Cure: Unraveling the Mystery of Hands-on Healing*. Louisville, CO: Sounds True, Inc; 2010.
13. Lindblad WJ, Diegelmann RF. Quantitation of hydroxyproline isomers in acid hydrolysates by high-performance liquid chromatography. *Anal Biochem*. 1984;138:390-395.
14. Ware JE, Jr, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care*. 1992;30:473-483.
15. McNair DM, Lorr M, Douglas MN, Fisher S, Droppelman LF. *Manual for the Profile Of Mood States*. San Diego, CA: Educational and Industrial Testing Service; 1971.
16. Rosenberg M. *Conceiving the Self*. Malabar, FL: Krieger; 1986.
17. Kurz A, Sessler DI, Lenhardt R. Perioperative normothermia to reduce the incidence of surgical-wound infection and shorten hospitalization. Study of wound infection and temperature group. *N Engl J Med*. 1996;334:1209-1215.
18. Wicke C, Halliday BJ, Scheuenstuhl H, Foree EF, Hunt TK. Examination of expanded polytetrafluoroethylene wound healing models. *Wound Repair Regen*. 1995;3:284-291.
19. Goodson WH 3rd, Hunt TK. Development of a new miniature method for the study of wound healing in human subjects. *J Surg Res*. 1982;33:394-401.
20. Radin DS, Stone J, Levine E, et al. Compassionate intention as a therapeutic intervention by partners of cancer patients: effects of a distant intention on the patients' autonomic nervous system, *Explore (NY)*; 2008;4:235-243.
21. Sloan RP, Ramakrishnan R. The MANTRA II study. *Lancet*. 2005;336:1769-1770; Author reply: 70-71.
22. Veber M. Why even a believer should not believe that god answers prayers, *Sophia* 2007; 46:177-187.
23. Hobbins PG. Compromised ethical principles in randomized clinical trials of distant, intercessory prayer. *J Bioethical Inq*. 2005;2:142-152.
24. Dossey L. *Be Careful What You Pray for. . . You Just Might Get It*. San Francisco, CA: HarperOne; 1998.
25. Fowler L. Praying for Obama's death. *Salon.com* 2009; Available at: [http://www.salon.com/2009/11/24/praying\\_for\\_obama\\_death/](http://www.salon.com/2009/11/24/praying_for_obama_death/). Accessed April 27, 2012.
26. Benson H, Dusek JA, Sherwood JB, et al. Study of the Therapeutic Effects of Intercessory Prayer (STEP) in cardiac bypass patients: a multicenter randomized trial of uncertainty and certainty of receiving intercessory prayer. *Am Heart J*. 2006;151:934-942.
27. Walker SR, Tonigan JS, Miller WR, Corner S, Kahlich L. Intercessory prayer in the treatment of alcohol abuse and dependence: a pilot investigation. *Altern Ther Health Med*. 1997;3:78-86.
28. Schmidt S, Schneider R, Utts J, Walach H. Distant intentionality and the feeling of being stared at: two meta-analyses. *Br J Psychol (Lond., England)*. 1953;2004:235-247.
29. Schlitz M, Braud W. Distant intentionality and healing: assessing the evidence. *Altern Ther Health Med*. 1997;3:62-73.
30. Walburn JV, Vedhara K, Hankins M, Rixon L, Weinman J. Psychological stress and wound healing in humans: A systematic review and meta-analysis. *J Psychosom Res*. 2009;67:253-271.